



Aerial Surveys of Distribution and Abundance of Marine Birds and Mammals in Santa Barbara Channel and the Santa Maria Basin

Final Technical Summary

Final Study Report



U.S. Department of the Interior
Minerals Management Service
Pacific OCS Region

Aerial Surveys of Distribution and Abundance of Marine Birds and Mammals in Santa Barbara Channel and the Santa Maria Basin

Final Technical Summary

Final Study Report

Authors

**Russell J. Schmitt
Michael L. Bonnell
Principal Investigators**

Prepared under MMS Cooperative
Agreement No. 14-35-0001-30758
by
Coastal Marine Institute
Marine Science Institute
University of California
Santa Barbara, CA 93106

U.S. Department of the Interior
Minerals Management Service
Pacific OCS Region

**Camarillo
March 2003**

Disclaimer

This report has been reviewed by the Pacific Outer Continental Shelf Region, Minerals Management Service, U.S. Department of the Interior and approved for publication. The opinions, findings, conclusions, or recommendations in this report are those of the authors, and do not necessarily reflect the views and policies of the Minerals Management Service. Mention of trade names or commercial products does not constitute an endorsement or recommendation for use. This report has not been edited for conformity with Minerals Management Service editorial standards.

Availability of Report

Extra copies of the report may be obtained from:

U.S. Dept. of the Interior
Minerals Management Service
Pacific OCS Region
770 Paseo Camarillo
Camarillo, CA 93010
Phone: 805-389-7621

A PDF file of this report is available at:
<http://www.coastalresearchcenter.ucsb.edu/CMI/>

Suggested Citation

The suggested citation for this report is:

Schmitt, Russell J. and Bonnell, Michael L. Aerial Surveys of Distribution and Abundance of Marine Birds and Mammals in Santa Barbara Channel and the Santa Maria Basin. MMS OCS Study 2003-012. Coastal Research Center, Marine Science Institute, University of California, Santa Barbara, California. MMS Cooperative Agreement Number 14-35-0001-30758. 23 pages.

Table of Contents

FINAL TECHNICAL SUMMARY	1
FINAL STUDY REPORT	3
Assessment of Historical Data for Sampling Design of the MMS Seabird Study	3
Introduction	4
Methods	6
Results and Discussion	6
Conclusions and Recommendations	14
Literature Cited	15
Seasonal Abundance and Distribution of Coastal Seabirds Offshore Santa Barbara and Ventura Counties, California	16
Abstract	16
Introduction	16
Materials and Methods	16
Results	17
Discussion	20
Acknowledgements	21
Literature Cited	21
Acknowledgements	23

Assessment of Historical Data for Sampling Design of the MMS Seabird Study

List of Tables

Table 1. Mean density of all alcids in Santa Maria Basin area, 1980-1983.	9
Table 2. Mean density of grebes, loons, scoters in Santa Maria Basin area, 1980-1983.	9
Table 3. Mean density of cormorants and pelicans in Santa Maria Basin area, 1980-1983.	10
Table 4. Number of sightings of prominent groups of marine birds in the Santa Maria Basin area, 1980-1983.	13

List of Figures

Figure 1. Existing Survey design for the MMS Seabird Study in the Santa Barbara Channel and the Santa Maria Basin.	5
Figure 2. Mean monthly density of alcids in the Santa Maria Basin area, 1980-1983.	6
Figure 3. Mean monthly density of grebes, loons, scoters in Santa Maria Basin area, 1980-1983.	8
Figure 4. Mean monthly density of cormorants and pelicans in Santa Maria Basin area, 1980-1983.	8
Figure 5. Mean monthly density, 1980-1983, of three groups of birds in the Santa Maria Basin area.	10
Figure 6 (a). Density (animals/km ²) of alcids for the months of July through October.	11
Figure 6 (b). Density (animals/km ²) of alcids for the months of January through March.	11
Figure 7. Density (animals/km ²) of loons, grebes, and scoters for the months of October through April.	12
Figure 8. Density (animals/km ²) of cormorants and pelicans for the months of October through April.	12

Seasonal Abundance and Distribution of Coastal Seabirds Offshore Santa Barbara and Ventura Counties, California

List of Tables

Table 1. MMS Coastal Seabird/Marine Mammal Aerial Surveys	17
Table 2. At-sea population estimates for coastal seabirds	18

List of Figures

Figure 1. Study area and standard aerial transect lines	17
Figure 2. Western grebe densities (number/km ²), September to March	18
Figure 3. Western grebe densities (number/km ²), April to August	18
Figure 4. Scoter densities (number/km ²), September to March	19
Figure 5. Scoter densities (number/km ²), April to August	19
Figure 6. Cormorant densities (number/km ²), all months	19
Figure 7. Loon densities (number/km ²), September to March	19
Figure 8. Loon densities (number/km ²), April to August	20
Figure 9. Brown pelican densities (number/km ²), February to May	20
Figure 10. Brown pelican densities (number/km ²), June to January	20

FINAL TECHNICAL SUMMARY

STUDY TITLE: Aerial Surveys of Distribution and Abundance of Marine Birds and Mammals in Santa Barbara Channel and the Santa Maria Basin

REPORT TITLE: Aerial Surveys of Distribution and Abundance of Marine Birds and Mammals in Santa Barbara Channel and the Santa Maria Basin

CONTRACT NUMBER: 14-35-0001-30758

SPONSORING OCS REGION: Pacific

APPLICABLE PLANNING AREA: Southern California

FISCAL YEARS OF PROJECT FUNDING: FY 96, FY 97

COMPLETION DATE OF THE REPORT: September 1998

COSTS: FY 96 - \$22,261, FY 97 – \$24,739

CUMULATIVE PROJECT COST: \$47,000

PROJECT MANAGER: Russell J. Schmitt

AFFILIATION: University of California, Santa Barbara

ADDRESS: Coastal Research Center, Marine Science Institute, University of California, Santa Barbara, CA 93106

PRINCIPAL INVESTIGATORS: ¹Russell J. Schmitt, ²Michael L. Bonnell (deceased)

ADDRESSES: ¹ Coastal Research Center, Marine Science Institute, University of California, Santa Barbara, CA 93106; ²Institute of Ocean Sciences, University of California, Santa Cruz, CA 95064

BACKGROUND: In November of 1994 an ambitious project was undertaken to provide detailed mapping of abundance and distribution of seabirds and marine mammals in the Santa Barbara Channel and the Santa Maria Basin (Pacific OCS Region). Our task was to separate the background noise from the significant patterns of use of the more than 100 species that occupy these waters. This series of bi-monthly aerial surveys was supported by the Marine Science Institute (MSI), UCSB, and the California Department of Fish and Game, Office of Spill Prevention and Response (OSPR). OSPR provided matching funds in the form of aircraft services and this work served their needs as well as the Minerals Management Service (MMS).

OBJECTIVES: Surveys continued through 1998 to gather additional data for geographical stratification of this area of development, abandonment, seismic profiling, and vessel traffic.

Stratification was based on an analysis of variance for the many species of birds that nest, roost, or migrate through this area. Regions with the highest variance received the greatest effort on surveys; regions with the lowest variance (i.e., most predictable patterns of use) received less effort.

DESCRIPTION AND RESULTS: Aerial Surveys were flown in a high-wing, twin-engine Partenavia PN68 Observer aircraft supplied by the CDFG Department of Air Services. The navigation was by GPS (Global Positioning System) with time/position entries into a computer at approximately eight-second intervals. The computer operator also entered important sightings (flocks of birds, marine mammals, water-mass boundaries, sea state, etc.) as well as number of observer(s) on watch. Typically only the observer on the shaded side of the aircraft was on duty; however, on northbound transects both observers were on watch searching a 50-meter corridor (a strip-census), with incidental sightings along the shore or outside the clinometer-defined search corridor noted. Sightings were recorded on voice-tape and later transcribed for computer entry by observation/time by the MMS. Time was then used to determine the position of sightings by melding the sighting file with the GPS record of time and latitude/longitude of the aircraft. All transects were flown at 200-foot (60-meters) altitude. To collect data on especially interesting sightings (e.g., flocks of birds near fishing boats, large whales, etc.), the aircraft increased altitude to approximately 600 ft.

From the beginning of this project, a total of 38 surveys were conducted (exclusive of commutes) in 130 hours in the air. Transects thus far include nearly 15,700 km (8,631 nautical miles) making this the most effort-intensive study conducted in any MMS-OCS Region. The prospects of this study are exciting; the last time systematic surveys were conducted in the Santa Barbara Channel and Santa Maria Basin was more than a decade ago.

SIGNIFICANT CONCLUSIONS: Over 50,000 sightings have been recorded, representing several hundred-thousand birds — this within a search swath of, at most, 100 meters. The numbers are astounding, in and of themselves, but we have also noted some remarkable findings — sightings of alcids (auklets and murrelets) appear sparse, numbers of gulls and pelicans seem stable, multiple observations of common dolphins are stable but expanding within their range, and large numbers of blue whales and humpback whales are clearly occupying the outer Santa Barbara Channel and Santa Maria Basin in mating/migration season.

STUDY PRODUCTS:

Bonnell, M. L. and R. G. Ford. 1996. Assessment of historical data for sampling design of the MMS seabird study. Prepared for Minerals Management Service, Pacific OCS Region, Camarillo, California, by Ecological Consulting, Inc., Portland, OR. 16 pp.

Pierson, Mark O., Michael D. McCrary, and Michael L. Bonnell. 1999. Seasonal Abundance and Distribution of Coastal Seabirds Offshore Santa Barbara and Ventura Counties, California. Fifth California Islands Symposium, OCS Study, MMS 99-0038: 428-434.

FINAL STUDY REPORT

ASSESSMENT OF HISTORICAL DATA FOR SAMPLING DESIGN OF THE MMS SEABIRD STUDY

Prepared for:

**Minerals Management Service
Pacific OCS Region**

Prepared by:

Michael L. Bonnell and R. Glenn Ford
Ecological Consulting, Inc.
2735 N.E. Weidler Street
Portland, OR 97232


October 1996

Introduction

The objective of this analysis was to use historical data to assess the adequacy of the aerial survey design presently used for the MMS Seabird Study. Available data were limited by coverage, both spatially and temporally, of previous systematic surveys conducted under contracts between the University of California, Santa Cruz (UCSC) and the Minerals Management Service, Pacific OCS Region. Surveys were conducted from 1975 through 1978 in the Southern California Bight and from 1980 through 1983 in waters of central and northern California. Methods are described in Briggs et al. 1981, 1983.

The regional focus of the MMS Seabird Study is waters of three coastal counties - Ventura, Santa Barbara, and San Luis Obispo. This area has been defined somewhat broadly in the first year of surveys to include all of the Santa Barbara Channel, nearshore waters along the mainland and north side of the Channel Islands, and the Santa Maria Basin offshore to about 121°00' W. longitude (Figure 1). Historical data on seabird use of these waters was limited to the two studies identified above; both attempted broad scale sampling of very large areas -- the entire Southern California Bight, and waters of central and northern California (Point Conception to Oregon). Data-collection methodology was nearly identical; however, studies substantially differed in survey effort. Surveys in the Santa Barbara Channel, 1975-1978, consisted of four transects, separated by 30' of longitude (about 48 km), flown north-south, plus a single east-west transect in mid-Channel. Aerial surveys were flown at approximately six-week intervals. In contrast, surveys in the Santa Maria Basin, 1980-1983, were conducted monthly and transect coverage was more than three times as great. For this reason, and the problematic difference in years of the two studies, we chose to examine the seasonal pattern of habitat use by seabirds in only the Santa Maria Basin area. If data were adequate for spatial stratification, we believed that these waters would provide the greatest chance of success.

Figure 1. Existing Survey design for the MMS Seabird Study in the Santa Barbara Channel and the Santa Maria Basin.



Methods

The limits of analysis were defined by waters from 34°30' to 35°30' N. latitude and extended as far offshore as survey transects. Using the Marine Mammal and Seabird Computer Database Analysis System (MMS-CDAS; ECI 1992), data were corrected for effort and binned into cells of a 5' grid of latitude and longitude for three species-groups: 1) alcids, 2) grebes, loons, and scoters, and 3) cormorants and pelicans. These groups were chosen on the basis of taxonomic relationship and habitat affinities. Initially, mean monthly densities were calculated for each species-group. The results were examined for significant interannual variation in species composition and area-wide abundance to identify the best survey windows (i.e., time of the year when greatest abundance was likely to occur). Without accessing individual records of observation conditions, no attempt could be made to exclude data recorded under less than ideal observation conditions. However, this was not viewed to be a problem because all data on these surveys were recorded by single observers on the glare-free side of the aircraft. We examined distributional patterns for alcids within the study area on a monthly basis, combining results for each month over all three years (see attached slide show). Based on this analysis, two seasons were selected: August and September, and January and February. The seasons were based on the variable distribution of alcids which show an offshore shift in the winter; other species-groups remain inshore throughout the year.

Results and Discussion

Sampling effort for this area was remarkably good -- few unsampled cells of a 5' grid were apparent in seasonal composites. Total area searched (1,046 sq. km) over three years was approximately 7% of the total area selected for analysis. Mean density of each of three species-groups in each month, February 1980 through January 1983, are shown in Figures 2-4 and detailed in Tables 1-3. Monthly density averaged over three years are shown for each species-group in Figure 5. Alcids were clearly most abundant in August and September, declining

through the late fall, followed by a significant peak in January. Maximum abundance of alcids in the summer was found close inshore; during winter, highest densities were reached farther from shore and beyond the area presently surveyed by the MMS Seabird Study (Figure 6). Numbers of grebes, loons, and scoters increased during the fall reaching a highly variable peak in the winter months (Figure 7). Cormorants were never very abundant in these waters, perhaps due to paucity of roosting sites and the relatively high level of human activities; greatest densities were observed in late fall. The inshore abundance of birds substantially increased in late summer and fall as Brown Pelicans migrated through the area (Figure 8).

Figure 2. Mean monthly density of alcids in the Santa Maria Basin area, 1980-1983.




Figure 3. Mean monthly density of grebes, loons, scoters in Santa Maria Basin area, 1980-1983.




Figure 4. Mean monthly density of cormorants and pelicans in Santa Maria Basin area, 1980-1983.




Table 1. Mean density of all alcids in Santa Maria Basin area, 1980-1983.

	1980	1981	1982	1983	ANNUAL MEAN	SD
Jan	-	7.48	7.89	8.18	7.850	0.352
Feb	2.23	6.03	1.47	-	3.243	2.443
Mar	2.31	0.67	0.84	-	1.273	0.902
Apr	0.34	0.36	0.34	-	0.347	0.012
May	0.77	4.78	0.58	-	2.043	2.372
Jun	1.24	1.22	0.74	-	1.067	0.283
Jul	2.81	4.98	0.71	-	2.833	2.135
Aug	23.73	1.47	0.51	-	8.570	13.138
Sep	17.15	19.76	0.68	-	12.530	10.345
Oct	8.27	5.84	1.12	-	5.077	3.636
Nov	4.70	1.47	3.21	-	3.127	1.617
Dec	0.83	3.95	2.70	-	2.493	1.570

Table 2. Mean density of grebes, loons, scoters in Santa Maria Basin area, 1980-1983.

	1980	1981	1982	1983	ANNUAL MEAN	SD
Jan	-	13.78	0.62	3.83	6.077	6.862
Feb	1.51	1.82	2.56	-	1.963	0.539
Mar	2.28	14.11	3.36	-	6.583	6.541
Apr	0.72	15.37	3.78	-	6.623	7.728
May	0.37	0.88	0.09	-	0.447	0.401
Jun	0.17	5.72	0.03	-	1.973	3.245
Jul	2.31	0.09	0.71	-	1.037	1.145
Aug	0.34	0.63	0.00	-	0.323	0.315
Sep	2.10	0.17	0.00	-	0.757	1.166
Oct	7.66	1.22	0.06	-	2.980	4.094
Nov	9.33	4.17	1.53	-	5.010	3.967
Dec	7.57	0.42	2.24	-	3.410	3.716

Table 3. Mean density of cormorants and pelicans in Santa Maria Basin area, 1980-1983.

	1980	1981	1982	1983	ANNUAL MEAN	SD
Jan	-	0.37	0.54	0.43	0.447	0.086
Feb	0.17	0.58	0.20	-	0.317	0.229
Mar	0.03	0.82	0.15	-	0.333	0.429
Apr	0.07	0.00	0.07	-	0.047	0.040
May	0.05	0.29	0.00	-	0.113	0.115
Jun	0.29	0.13	0.03	-	0.150	0.131
Jul	2.39	1.22	0.08	-	1.230	1.155
Aug	1.24	0.11	0.06	-	0.470	0.667
Sep	1.33	0.11	0.12	-	0.520	0.701
Oct	6.88	0.30	0.26	-	2.480	3.811
Nov	10.99	0.30	0.06	-	3.783	6.242
Dec	0.26	0.73	0.16	-	0.383	0.304

Figure 5. Mean monthly density, 1980-1983, of three groups of birds in the Santa Maria Basin area.




Figure 6 (a). Density (animals/km²) of alcids for the months of July through October, averaged over the three years of available data. Data were binned into 15 minute blocks for contouring.




Figure 6 (b). Density (animals/km²) of alcids for the months of January through March, averaged over the three years of available data. Data were binned into 15 minute blocks for contouring.




Figure 7. Density (animals/km²) of loons, grebes, and scoters for the months of October through April, averaged over the three years of available data. Data were binned into 15 minute blocks for contouring.





Figure 8. Density (animals/km²) of cormorants and pelicans for the months of October through April, averaged over the three years of available data. Data were binned into 15 minute blocks for contouring.



Some indication of the seasonal change in faunal composition can be obtained simply on the basis of number of sightings (Table 4). Auklets were the most abundant of the alcids in all months except July through September; during those months, Common Murres predominated. On a yearly average, Cassin's Auklet accounted for 48% of all alcids and 74% of all auklets. Rhinoceros Auklets were rare or absent from April through October. Murrelets were infrequently identified but may have been recorded as unidentified small alcids. Sightings of grebes, loons, and scoters in inshore waters of the Santa Maria Basin were nearly equal within groups. The only grebe identified to species level was the Western Grebe; the unidentified grebes category represented only about 1.5% of total grebes. Peak numbers of Western Grebes were recorded from about October through March. On a yearly average, Pacific loons accounted for about 74% of all loons; Red-throated Loons were present in small numbers from January through April and, interestingly, Common Loons were identified only in April and May, and from August through November. Both Surf and White-winged Scoters were identified to the species level, and Surf Scoters predominated in all months except January. It should be noted that sightings of scoters identified to the species level represented only 54% of all records of scoters. Cormorants were recorded in all months with greatest numbers from November through February. Brown Pelicans accounted for 44% of all sightings in this group and, although present in small numbers year round, only reached substantial abundance from August through December.

Table 4. Number of sightings of prominent groups of marine birds in the Santa Maria Basin area, 1980-1983.

	Murres	Auklets	Unid. Alcids	Grebes	Loons	Scoters	Corm. Sp.	Brown Pelican
Jan	125	217	61	30	18	34	10	2
Feb	26	162	53	16	31	17	20	2
Mar	5	39	35	15	20	23	5	2
Apr	3	16	8	4	40	16	3	0
May	8	43	13	7	8	0	4	1
Jun	2	27	17	2	3	1	4	1
Jul	23	15	5	0	2	3	1	8
Aug	55	7	7	3	2	4	5	9
Sep	70	12	11	5	1	0	5	11
Oct	63	61	8	24	2	6	8	16
Nov	15	76	84	23	28	14	22	10
Dec	18	96	29	6	8	31	6	11

Conclusions and Recommendations

Timing of surveys. Through the first year of the MMS Seabird Study, and proposed to the State's Office of Aircraft Services for subsequent work in 1996 and 1997, surveys would be flown according to the following schedule:

January
February
April
May
June
August
October
November

Some limitations in scheduling result from other commitments of the aircraft (including other surveys conducted by UCSC on behalf of the Office of Oil Spill Prevention and Response, OSPR, in other waters of California). The original schedule of surveys was made in consultation with K. T. Briggs, and the months appear to be well chosen. On the basis of our examination of historical data, no changes in the schedule are recommended.

Spatial coverage and survey effort. To a great extent, the level of sampling effort that can be devoted to the Santa Maria Basin area is determined by hours of aircraft availability and day length in the winter months. The present set of transects (Figure 1) provide thorough coverage of both the Santa Barbara Channel and the Santa Maria Basin within existing logistical constraints. The results of our analysis do not suggest that changes be made in the present study design. However, additional analysis should be done using the database produced by the first year of the MMS Seabird Study to identify areas that should receive greater survey effort.

Literature Cited

Briggs, K. T., E. W. Chu, D. B. Lewis, W. B. Tyler, R. L. Pitman, and G. L. Hunt. 1981. Distribution, Numbers, and Seasonal Status of Seabirds of the Southern California Bight. Part III of Investigator's Reports, Summary of Marine Mammal and Seabird Surveys of the Southern California Bight Area, 1975-1978. Final report prepared by the Univ. of Calif., Santa Cruz, for the Bureau of Land Management, U. S. Dept. of the Interior. National Technical Information Service # PB-81-248-197, Springfield, VA.

Briggs, K. T., W. B. Tyler, D. B. Lewis, and K. F. Dettman. 1983. Seabirds of central and northern California, 1980-1983: Status, abundance, and distribution. Investigator's Final Report, prepared by the Univ. of Calif., Santa Cruz, for the Minerals Management Service, Pacific OCS Region. OCS Study MMS 84-0043.

Ecological Consulting, Inc. 1992. User's Guide and Database Documentation: Marine mammal and seabird computer database analysis system. Prepared for the Minerals Management Service, Pacific OCS Region, by Ecological Consulting, Inc. Portland, OR.

SEASONAL ABUNDANCE AND DISTRIBUTION OF COASTAL SEABIRDS OFFSHORE SANTA BARBARA AND VENTURA COUNTIES, CALIFORNIA

Mark O. Pierson¹, Michael D. McCrary¹, and Michael L. Bonnell²

¹Pacific OCS Region, Minerals Management Service, 770 Paseo Camarillo, Camarillo, CA 93010
(805) 389-7863, FAX 389-7874, E-mail: mark.pierson@mms.gov, michael.mccrary@mms.gov

²Institute of Marine Sciences, University of California, Santa Cruz, CA 95064
(831) 426-5844, E-mail: eci@netcom.com

ABSTRACT

Between January 1996 and August 1997, the Minerals Management Service (MMS), in cooperation with the University of California, Santa Cruz (UCSC), and the California Department of Fish and Game's Office of Oil Spill Prevention and Response (OSPR), conducted 12 aerial surveys of seabirds and marine mammals in the waters of the Santa Barbara Channel and the Santa Maria Basin. The goals of the study were to identify and characterize important habitats of seabirds known to be especially sensitive to oil spill impacts, and to update and provide a database for comparison with earlier studies of marine mammals and seabirds in these waters. This paper focuses on one aspect of this ongoing study: aerial surveys of divers and brown pelicans in the nearshore waters along the mainland shoreline of Santa Barbara and Ventura counties and along the north side of the northern Channel Islands. Western/Clark's grebes were the most abundant nearshore seabirds during most months of the year, reaching densities as high as 400 birds/km² in some areas. Other abundant species or groups of species included scoters (primarily surf scoters), brown pelicans, cormorants, and loons. Highest seabird densities consistently were recorded near Point Conception and along the Ventura County coast.

Keywords: Abundance, distribution, nearshore waters, seabirds, seasonality, southern California, western grebe (*Aechmophorus occidentalis*), Clark's grebe (*Aechmophorus clarkii*).

INTRODUCTION

In January 1996, the Minerals Management Service (MMS), in cooperation with the University of California, Santa Cruz (UCSC), and the California Department of Fish and Game's Office of Oil Spill Prevention and Response (OSPR), began a study of the abundance and distribution of seabirds and marine mammals in the Santa Barbara Channel and Santa Maria Basin. The last systematic studies of seabird distribution in these waters had been conducted from 1975 through 1983 (Briggs et al. 1981, 1983). These studies were large-scale, relatively coarsely grained surveys, which did not focus on nearshore seabird populations. Additionally, in the years since those studies many changes

occurred in oceanographic conditions and the breeding status of colonial seabirds (Sowls et al. 1980; Carter et al. 1992).

The goals of the current MMS study were to identify and characterize important habitats of marine birds known to be especially sensitive to oil spill impacts, and to update and provide a database for comparison with the earlier studies of marine birds in these waters. The principal areas of interest were 1) nearshore populations of divers, mostly migrant and overwintering species in southern California (loons, grebes, cormorants, and scoters); and 2) pelagic populations, which include both migrants and local breeders (alcids, shearwaters, phalaropes, gulls, and pelicans). The coastal divers and the more pelagic alcid species, such as murres and auklets, are considered to be particularly vulnerable to the effects of oiling.

The information on seabird and marine mammal distribution and abundance collected during this study has been added to the MMS Pacific Outer Continental Shelf (OCS) Region's existing database (Ecological Consulting, Inc. 1992). These data can be used to identify important offshore habitats, be compared with previously acquired data to detect long-term changes in the distribution and abundance of key species, and be used to estimate potential impacts to these marine resources from accidental oil spills. This paper is a preliminary report on the results of the first 12 aerial surveys of the study, focusing on coastal seabird populations inhabiting the nearshore waters along the southern and central California mainland and the north side of the northern Channel Islands. More detailed analysis of the data will follow completion of the aerial surveys and compilation of the final study database.

MATERIALS AND METHODS

The California State Office of Aircraft Services, under support from OSPR, provided the survey aircraft and an experienced warden-pilot. The survey aircraft, a Partenavia PN68 Observer, was a two-engine, high-winged airplane that was well suited for overwater operations and equipped with bubble windows for observations.

The approach and methodology were first developed by UCSC biologists for Bureau of Land Management-funded baseline studies of marine mammals and seabirds offshore

southern California in the middle and late 1970s. The methodology was refined during subsequent MMS-funded baseline studies offshore central and northern California and off Oregon and Washington in the 1980s (Briggs et al. 1981, 1983, 1991).

On board the survey aircraft, a Global Positioning System (GPS) unit linked to a laptop computer was used to record latitude, longitude, and time at 5-sec intervals. Surveys were flown at an altitude of 60 m (200 ft) above sea level and ground speed of 90 kts. Observers searched a 50-m corridor on each side of the aircraft; boundaries were strictly defined using a clinometer and simple trigonometry. Both sides of the aircraft trackline were worked under optimum conditions. The 60-m altitude allowed the smallest birds (e.g., auklets, murrelets, storm-petrels, and phalaropes) to be seen and usually identified to the species level under good light conditions (Briggs et al. 1985a, b). Observations were recorded on audio-cassette and included counts; species; behavior; associations with boats, plankton, water-mass boundaries, other birds, mammals, etc.; and the time.

Post-survey computer data entry was done from transcriptions of audio-cassette tapes to produce a database of sighting information. For each observation, latitude and longitude were estimated using the time for interpolation between the previous and subsequent position fixes. These data were used to create GIS data layers for the aircraft trackline and for locations of bird observations. Numbers of birds observed and area sampled were binned into 5°-latitude/longitude blocks for the purpose of calculating densities. Densities, represented as birds/km², were displayed as rectangular latitude/longitude blocks and as isopleths of density. Density isopleths were generated using a standard linear interpolation model and smoothed with a Bezier spline.

The survey design was developed to optimize the sampling scheme, while balancing the level of areal coverage and the available airtime. Survey tracklines were flown in the two principal areas of interest: nearshore, along the mainland and the north side of the Channel Islands; and offshore in the Santa Barbara Channel and the Santa Maria Basin west to about 121°W longitude (Figure 1). The nearshore mainland tracklines extended from Leo Carrillo State Beach west and north to the Santa Maria River mouth (encompassing the Ventura and Santa Barbara County shorelines). The nearshore island tracklines extended from the east end of Anacapa Island west to Point Bennett on San Miguel Island. Based on analysis of historical data on seasonality and distribution, the months chosen for survey were January, February, April, May, June, August, October, and November (Bonnell and Ford 1996).

The California Marine Mammal and Seabird Database (Ecological Consulting, Inc. 1992), which incorporates sighting data from all previous California baseline surveys, made it possible to compare data collected offshore southern California over several decades and track possible changes in species composition, numbers, and distribution over time.

RESULTS

In this paper, we report the results of 12 nearshore aerial surveys conducted along the mainland and the north side of the northern Channel Islands, from January 1996 to August 1997 (Figure 1; Table 1). The paper focuses on the most abundant nearshore species of divers and the brown pelican (*Pelecanus occidentalis*). Among the divers, western/Clark's grebes (*Aechmophorus occidentalis/clarkii*) were the only species selected for analysis. The loons, cormorants, and scoters, which are all more difficult to identify to the species level from the air, were each grouped for analysis.

Although several species of gulls and, to a lesser extent, terns also were abundant and important components of the nearshore avifauna, they are not discussed in this paper. The same is true of alcids, phalaropes, and shearwaters, even though the latter occasionally occupied localized nearshore areas in the tens of thousands. These species will be considered in future works, which will include data from both nearshore and open-water transects.



Figure 1. Study area and standard aerial transect lines.

Table 1. MMS Coastal Seabird/Marine Mammal Aerial Surveys included for analysis in this paper.

Survey Number	Dates
96-1	22-23 January 1996
96-2	22-23 February 1996
96-3	9-10 April 1996
96-4	8-9 May 1996
96-5	26-27 June 1996
96-6	12-13 August 1996
96-7	24-25 October 1996
96-8	11 November 1996
97-1	26-27 February 1997
97-2	9-10 April 1997
97-3	11-12 June 1997
97-4	13-14 August 1997

Pierson, M. O., et al.

Western/Clark's Grebes

These two species could not be distinguished from the air and henceforth will be referred to simply as 'western' grebes. Densities of western grebes were calculated for two periods: September through March, which encompasses most of the fall and spring migrations, plus the overwintering birds; and April through August, when most adults are nesting at freshwater locations in the western U.S. and Canada (Palmer 1962; Garrett and Dunn 1981).

During the September -to-March period, highest densities of western grebes were recorded in nearshore waters along the Ventura County shoreline and Santa Barbara County north of Point Conception, particularly between Arguello and Purisima points (Figure 2). Sampled densities reached a peak of 400 birds/km² off western Ventura County. Based on the recorded densities, we estimated that approximately 71,300 western grebes were present at sea in the study area during this period (Table 2).

Recorded densities were substantially lower during the April-to-August period (Figure 3). Although relatively high numbers of grebes may still be present in area waters in April, near the end of the spring migration, numbers observed dropped sharply during the summer months. Only a few hundred western grebes were recorded during the August 1996




Figure 2. Western grebe (*Aechmophorus occidentalis/clarkii*) densities (number/km²) recorded for the months of September through March.

Table 2. At-sea population estimates for coastal seabirds derived from data collected on MMS Seabird Study aerial surveys from January 1996 through August 1997.

Seabird	Survey Months	Density/km ²		Surveyed Area (km ²)	Population Estimate
		Mean	Standard Deviation		
Western Grebe	Sep-Mar	13.58	43.45	8,902.2	71,320
	Apr-Aug	3.66	15.77	9,258.8	21,591
Scoters	Sep-Mar	6.41	27.45	8,901.1	35,696
	Apr-Aug	1.37	5.71	9,258.8	8,965
Cormorants	Jan-Dec	1.49	4.27	9,542.4	10,455
Loons	Sep-Mar	0.79	4.13	8,899.7	5,971
	Apr-Aug	0.39	1.05	9,258.2	2,562
Brown Pelican	Feb-May	0.95	2.32	9,327.4	8,342
	Jan-Jan	1.38	3.80	9,186.7	10,363

aerial survey. As was the case for the September to March period, highest densities were recorded along the Ventura County shoreline and north of Point Conception. Sampled densities reached a peak of 142 birds/km² off western Ventura County. Based on the recorded densities, we estimated that approximately 21,600 western grebes were present at sea in the study area during this period (Table 2).




Figure 3. Western grebe (*Aechmophorus occidentalis/clarkii*) densities (number/km²) recorded for the months of April through August.

Scoters

Three species of scoters occur in southern and central California waters: surf scoter (*Melanitta perspicillata*), white-winged scoter (*M. fusca*), and black scoter (*M. nigra*). The vast majority of scoters in the study area waters (and of those identified from the air) are surf scoters; black scoters are relatively uncommon (Garrett and Dunn 1981; Briggs et al. 1987).

As was the case for western grebes, densities of scoters were calculated for the September-to-March and April-to-August periods. The highest overall densities during September to March were observed at the Channel Islands and along the mainland north of Point Conception (Figure 4). Peak densities of nearly 267 birds/km² were recorded north of Point Sal near the Santa Maria River mouth. Based on the recorded densities, we estimated that approximately 35,700 scoters were present at sea in the study area during this period (Table 2).

During April to August, overall scoter densities declined (Figure 5). Highest densities again were recorded along the mainland north of Point Conception, while moderate densities were observed off Ventura County and at the Channel Islands. Peak densities of 51 birds/km² were recorded in nearshore waters north of Purisima Point. Based on the recorded densities, we estimated that approximately 9,000 scoters were present at sea in the study area during this period (Table 2).

Cormorants

The three cormorant species occurring in the study area include Brandt's cormorant (*Phalacrocorax*




Figure 4. Scoter (*Melanitta* spp.) densities (number/km²) recorded for the months of September through March.




Figure 6. Cormorant (*Phalacrocorax* spp.) densities (number/km²) recorded for all months surveyed.




Figure 5. Scoter (*Melanitta* spp.) densities (number/km²) recorded for the months of April through August.

scattered along the mainland and Channel Island shorelines (Figure 7). The peak densities of about 9 birds/km² were recorded at Santa Rosa Island. Based on the recorded densities, we estimated that approximately 6,000 loons were present at sea in the study area during this period (Table 2).
 The numbers of loons in the study area dropped during April to August, when highest densities were recorded along the mainland north of Point Conception, at the east end of the Santa Barbara Channel, and at the Channel Islands (Figure 8). Peak densities of approximately 9 birds/km² were observed along the mainland north of Purisima Point. Based on the recorded densities, we estimated that approximately 2,600 loons were present at sea in the study area during this period (Table 2).

penicillatus), double-crested cormorant (*P. auritus*), and pelagic cormorant (*P. pelagicus*). All three species breed locally. Since the cormorant species could not always be reliably distinguished from the air, they were combined for analysis.

Because cormorants are year-round residents in the study area, densities were calculated for all months surveyed (Figure 6). Cormorants were most abundant in nearshore waters, especially along the southwestern Santa Barbara County shoreline and around the Channel Islands. Peak densities of approximately 49 birds/km² were observed along the mainland near Gaviota. Based on the recorded densities, we estimated that approximately 10,500 cormorants were present at sea in the study area (Table 2).

Loons

Three species of loon migrate through and overwinter in southern California waters: Pacific loon (*Gavia pacifica*), red-throated loon (*G. stellata*), and common loon (*G. immer*). Because the loon species could not always be reliably distinguished from the air, they were combined for analysis.

Loon densities were calculated for the September-to-March and April-to-August periods. During September to March, relatively low densities of loons were observed




Figure 7. Loon (*Gavia* spp.) densities (number/km²) recorded for the months of September through March.

Brown Pelican

The brown pelican, which currently breeds in our study area only on Anacapa Island, is listed as an endangered species under the federal Endangered Species Act of 1973. On aerial surveys conducted during the months of February through May, corresponding to the pelican breeding season, densities of up to 26 birds/km² were recorded (Figure 9). Highest densities were observed near the rookery on West

Pierson, M. O., et al.



Figure 8. Loon (*Gavia* spp.) densities (number/km²) recorded for the months of April through August.

Anacapa Island and along the mainland in the eastern Santa Barbara Channel. Based on the recorded densities, we estimated that approximately 8,200 brown pelicans were present at sea in the study area during this period (Table 2).

Higher overall densities were recorded outside the breeding season, from June through January, when birds from the larger Mexican colonies moved northward into southern California (Figure 10). Brown pelicans were somewhat more widely distributed than during the breeding season, with highest densities recorded along the mainland shoreline from Point Arguello eastward and near the Channel Islands. Peak densities of about 48 birds/km² were observed near Point Conception. Based on the recorded densities, we estimated that approximately 10,400 brown pelicans were present at sea in the study area during this period (Table 2).

DISCUSSION

The results reported in this paper are preliminary, since they represent only the first 12 aerial surveys of the MMS Seabird Study. However, some generalizations can be drawn about the seasonal distribution and abundance of the selected nearshore species. Additionally, some limited comparisons can be made with earlier seabird research in the study area.

Western grebes were the most abundant coastal seabirds in the study area except during the late spring and summer months. Bonnell and Ford (1996), reanalyzing data from earlier MMS studies in our study area (Briggs et al. 1981, 1983), reported that peak numbers of western grebes were recorded from about October through March. Similarly, Briggs et al. (1987), in their monograph on California seabird communities, identified western grebes as one of the predominant nearshore species along the mainland from October to May. They estimated a peak winter population of about 27,000 birds south of Point Conception, with most birds concentrated in shallow waters at the eastern end of the Santa Barbara Channel. Summer populations were much smaller, on the order of 500 to 800 birds.

Although we estimate a much larger fall-winter population of western grebes (approximately 71,300) than that presented by Briggs et al. (1987), the two estimates cannot



Cormorants were the third most abundant group of nearshore divers in our study area, with a mean at-sea population estimate of approximately 10,400 birds. Briggs et al. (1987) found Brandt's cormorants to be the most abundant of the three cormorant species in southern California, with highest densities recorded from late summer through late spring. They estimated that numbers in southern California increased from about 5,500 nesting birds in late spring and summer to more than 20,000 by late September, with the arrival of central California birds after the nesting season. Carter et al. (1992) estimated that more than 27,000 cormorants bred in our study area in 1989 to 1991. Almost all of these were on the northern Channel Islands, and more than 85% were Brandt's cormorants.

Briggs et al. (1987) found Pacific loons to be most abundant in southern California waters during fall migration, estimating that 40,000 to 60,000 may have been present at one time in mid-December. Bonnell and Ford (1996) reported that Pacific loons accounted for about 74% of all loons identified during the earlier MMS studies (Briggs et al. 1981, 1983). Although we did not observe such abundance, the timing of our surveys (i.e., we surveyed the study area only once in the months of November and December) made it difficult to observe such brief migratory events (Table 1). One of us (Pierson, unpubl. obs.) observed thousands of loons migrating southward across the western Santa Barbara Channel in late November 1995.

According to Briggs et al. (1987), only about 5,000 Pacific loons overwintered in California waters south of Point Conception in each of the winters from 1976 to 1978, although densities as high as 80 birds/km² were recorded over the shallows east of Anacapa Island. In nearshore mainland waters, red-throated loons were more abundant, with estimated peak numbers of 1,000 to 3,000 birds. Common loons were recorded in greatest numbers north of Point Conception, where 5,000 to 10,000 birds were estimated to occur at sea during spring migration (March to May). Our estimate for the September-through-March period (approximately 6,000 birds) is generally comparable to the earlier figures for overwintering birds of all three species.

The brown pelican population continues to recover from the declines recorded in the 1960s and 1970s. Carter et al. (1992) estimated that nearly 12,000 brown pelicans nested in southern California in 1989 to 1991, most of these on West Anacapa Island in our study area. In general, our observations correspond well with those of the earlier studies (Briggs et al. 1981, 1983). Briggs et al. (1987) reported that pelican numbers were lowest from December through March, when 5,000 to 6,000 pelicans (half of which were breeders) were present in southern California. From about May through December, these numbers were augmented by tens of thousands of pelicans that moved north from the Mexican colonies. In September and October, when maximum numbers of pelicans were in California waters, 70 to 80% of the statewide population occurred south of Point Conception (resulting in an estimated population as high as 80,000 to 90,000 in southern California). Although our

population estimates are much lower (approximately 10,400), we surveyed only a portion of the species' southern California range and did not include pelicans onshore.

Although our results are preliminary, we can draw some conclusions about nearshore seabird populations in the study area. Overall, abundance was greatest in the fall and winter, between September and March or April. Greatest densities occurred along the Ventura County mainland shoreline (particularly between Mugu Lagoon and Rincon Point), and along the Santa Barbara County shoreline north of Point Conception (particularly between Point Arguello and the Santa Maria River mouth). Relatively high densities also were observed along the north sides of the northern Channel Islands, particularly in the vicinity of Anacapa and Santa Cruz islands.

Tens of thousands of seabirds occupied the nearshore waters during these months. Most of these birds were western grebes and surf scoters, although thousands of cormorants, loons, pelicans, and gulls were also present. Occasionally, aggregations of thousands to tens of thousands of shearwaters or phalaropes appeared in nearshore waters along the mainland or islands.

Numbers were much lower during the summer months, when the nearshore avifauna was dominated by locally breeding gulls and cormorants and by brown pelicans. Overall distribution patterns during these months did not differ substantially from those observed during fall and winter.

Completion of the MMS Seabird Study database will allow us to address these and other questions on the abundance and distribution of seabirds in the study area. We also hope to be able shed some light on the possible effects of the 1997 to 1998 El Niño episode on area seabird populations.

ACKNOWLEDGMENTS

We would like to thank a number of people for their help during the course of this study and the preparation of this paper. Ken Briggs, Breck Tyler, Dave Lewis, and Jeff Davis all lent their experience and expertise to the field effort, serving as observers on numerous surveys. Our primary pilots, Bob Morgan and Larry Heit, flew our surveys with great skill and patience, often under difficult circumstances. Glenn Ford developed and upgraded the database and GIS programs used for this study and produced the density and population estimates for this paper. Finally, Ken Briggs, Glenn Ford, and Cathie Dunkel reviewed the early drafts of this paper and provided many valuable suggestions.

LITERATURE CITED

- Bonnell, M. L. and R. G. Ford. 1996. Assessment of historical data for sampling design of the MMS seabird study. Prepared for Minerals Management Service, Pacific OCS Region, Camarillo, California, by Ecological Consulting, Inc., Portland, OR. 16 pp.

Pierson, M. O., et al.

- Briggs, K. T., E. W. Chu, D. B. Lewis, W. B. Tyler, R. L. Pitman, and G. L. Hunt, Jr. 1981. Distribution, numbers, and seasonal status of seabirds of the Southern California Bight. Book I, Part III. Investigators' Reports, summary of marine mammal and seabird surveys of the Southern California Bight area, 1975-1978. Publ. #PB-81-248-205, U.S. National Technical Information Service, Springfield, VA. 399 pp.
- Briggs, K. T., W. B. Tyler, and D. B. Lewis. 1985a. Comparison of ship and aerial surveys of birds at sea. *Journal of Wildlife Management* 49(2):405-411.
- Briggs, K. T., W. B. Tyler, and D. B. Lewis. 1985b. Aerial surveys for seabirds: methodological experiments. *Journal of Wildlife Management* 49(2):412-417.
- Briggs, K. T., W. B. Tyler, D. B. Lewis, and D. R. Carlson. 1987. Bird communities at sea off California: 1975 to 1983. Cooper Ornithological Society, Studies in Avian Biology No. 11. 74 pp.
- Briggs, K. T., W. B. Tyler, D. B. Lewis, and K. F. Dettman. 1983. Seabirds of central and northern California, 1980-1983: status, abundance, and distribution. Center for Marine Studies, University of California, Santa Cruz. Prepared for U.S. Department of the Interior, Minerals Management Service, Pacific OCS Region. Contract #AA551-CT9-33. 246 pp.
- Briggs, K. T., D. H. Varoujean, W. W. Williams, R. G. Ford, M. L. Bonnell, and J. L. Casey. 1991. Seabirds of the Oregon and Washington OCS, 1989-1990. Chapter III in Brueggeman, J. J. (ed.), Oregon and Washington marine mammal and seabird surveys. Draft Final Report prepared by Ebasco Environmental, Bellevue, Washington, and Ecological Consulting, Inc., Portland, Oregon, for the U.S. Department of the Interior, Minerals Management Service, Pacific OCS Region, Camarillo, CA. OCS Study MMS 91-000.
- Carter, H. R., G. J. McChesney, D. L. Jaques, C. S. Strong, M. W. Parker, J. E. Takekawa, D. L. Jory, and D. L. Whitworth. 1992. Breeding populations of seabirds in California, 1989-1991. U.S. Fish and Wildlife Service, Northern Prairie Wildlife Research Center, Dixon, California, and San Francisco Bay National Wildlife Refuge Complex, Newark, California. Draft final report to Minerals Management Service, Pacific OCS Region, under Inter-agency Agreement No. 14-12-001-30456. Volumes I and II.
- Ecological Consulting, Inc. 1992. Marine Mammal and Seabird Computer Database Analysis System. User's guide and database documentation, California Marine Mammal and Seabird Databases. Prepared for U.S. Department of the Interior, Minerals Management Service, Pacific OCS Region.
- Garrett, K. and J. Dunn. 1981. Birds of Southern California: Status and Distribution. Los Angeles Audubon Society, Los Angeles, CA. 408 pp.
- Palmer, R. S. 1962. Handbook of North American Birds. Volume 1: Loons through Flamingos. Yale University Press, New Haven and London. 567 pp.
- Sowls, A. L., A. R. DeGange, J. W. Nelson, and G. S. Lester. 1980. Catalog of California Seabird Colonies. Department of the Interior, U.S. Fish and Wildlife Service. FWS/OBS-80/37. 371 pp.

ACKNOWLEDGEMENTS

Michael McCrary and Dr. Mark Pierson were the major observers on most surveys and deserve a sincere debt of gratitude, as does the MSI at UCSB, Bonnie Williamson for her able administration, Ed Cassano for a permit to work within nearshore waters of the Channel Islands National Marine Sanctuary, and OSPR for their continued support. It is not often that such a long-term project receives such sustained funding. My thanks to all.



The Department of the Interior Mission

As the Nation's principal conservation agency, the Department of the Interior has responsibility for most of our nationally owned public lands and natural resources. This includes fostering sound use of our land and water resources; protecting our fish, wildlife, and biological diversity; preserving the environmental and cultural values of our national parks and historical places; and providing for the enjoyment of life through outdoor recreation. The Department assesses our energy and mineral resources and works to ensure that their development is in the best interests of all our people by encouraging stewardship and citizen participation in their care. The Department also has a major responsibility for American Indian reservation communities and for people who live in island territories under U.S. administration.



The Minerals Management Service Mission

As a bureau of the Department of the Interior, the Minerals Management Service's (MMS) primary responsibilities are to manage the mineral resources located on the Nation's Outer Continental Shelf (OCS), collect revenue from the Federal OCS and onshore Federal and Indian lands, and distribute those revenues.

Moreover, in working to meet its responsibilities, the **Offshore Minerals Management Program** administers the OCS competitive leasing program and oversees the safe and environmentally sound exploration and production of our Nation's offshore natural gas, oil and other mineral resources. The **MMS Royalty Management Program** meets its responsibilities by ensuring the efficient, timely and accurate collection and disbursement of revenue from mineral leasing and production due to Indian tribes and allottees, States and the U.S. Treasury.

The MMS strives to fulfill its responsibilities through the general guiding principles of: (1) being responsive to the public's concerns and interests by maintaining a dialogue with all potentially affected parties and (2) carrying out its programs with an emphasis on working to enhance the quality of life for all Americans by lending MMS assistance and expertise to economic development and environmental protection.